

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 27/00, 27/26, 33/543, 33/551, C12Q 1/68, A01N 1/02, C12M 1/00		A1	(11) International Publication Number: WO 98/08083 (43) International Publication Date: 26 February 1998 (26.02.98)
(21) International Application Number: PCT/US97/14372 (22) International Filing Date: 13 August 1997 (13.08.97) (30) Priority Data: 08/700,182 20 August 1996 (20.08.96) US (71) Applicant (for all designated States except US): MOTOROLA INC. [US/US]; 1303 East Algonquin Road, Schaumburg, IL 60196 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): REBER, William, L. [US/US]; 1029 Buccaneer Road #6, Schaumburg, IL 60916 (US). ACKLEY, Donald, E. [US/US]; 2033 Cambridge Avenue, Cardiff, CA 92007 (US). PERTTUNEN, Cary, D. [US/US]; 11764 Raintree Court, Shelby Township, MI 48315 (US). (74) Agents: SARLI, Anthony, J., Jr. et al.; Motorola Inc., Intellectual Property Dept., 1303 East Algonquin Road, Schaumburg, IL 60196 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: **METHOD AND APPARATUS FOR DETECTING PREDETERMINED MOLECULAR STRUCTURES IN A SAMPLE**

(57) Abstract

A predetermined molecular structure in a sample is detected by sensing a pattern in which the sample binds to an array of binding sites in a molecular detection device (20), and comparing the pattern to a reference pattern to detect the predetermined molecular structure in the sample (22). In one embodiment, the reference pattern is generated by sensing a pattern in which a reference sample containing the predetermined molecular structure binds to a like array of binding sites. In another embodiment, the reference pattern is generated by predicting a pattern in which the predetermined molecular structure binds to the array of binding sites.